

QCD on Coarse Lattices*

M. Alford, W. Dimm and G.P. Lepage

Floyd R. Newman Laboratory of Nuclear Studies, Cornell University, Ithaca, NY 14853

G. Hockney and P.B. Mackenzie

Theoretical Physics Department, Fermilab, Batavia, IL 60510

We show that the perturbatively-improved gluon action for QCD, once it is tadpole-improved, gives accurate results even with lattice spacings as large as 0.4 fm. *No* tuning of the couplings is required. Using this action and lattice spacing, we obtain a static potential that is rotationally invariant to within a few percent, the spin-averaged charmonium spectrum accurate to within 30–40 MeV, and scaling to within 5–10%. We demonstrate that simulations on coarse lattices are several orders of magnitude less costly than simulations using current methods.

* Talk presented by G.P. Lepage at *Lattice '94* (Bielefeld, Germany, September 1994)

Recent studies indicate that weak-coupling perturbation theory, when properly renormalized and tadpole improved, gives an accurate account of lattice-QCD dynamics even at distances as large as 1/2 fm [1]. This suggests that accurate simulations can be carried out on very coarse lattices, perhaps with lattice spacing $a \approx 1/2$ fm, by using (asymptotically free) perturbation theory to correct for dynamics at distance scales smaller than a . In this paper we present new evidence that this is indeed the case for gluon dynamics. We show accurate results obtained with spacing $a \approx 0.4$ fm. This is 4–8 times larger than usual a 's, and leads to a spectacular reduction in simulation costs, which typically vary as $1/a^6$.

The standard discretizations of the quark and gluon actions are not sufficiently accurate for simulations on such coarse lattices; the leading finite- a errors must be removed. Such errors can be systematically removed from on-shell quantities, like scattering amplitudes and masses, by adding nonrenormalizable terms to the lattice actions [2]. Since these terms correct primarily the short-distance behavior of the theory, their coefficients can be computed using perturbation theory once a is sufficiently small. Early tests of this technique for lattice QCD showed little improvement, but these used naive lattice perturbation

theory and so seriously underestimated the coefficients. The use of renormalized perturbation theory and especially tadpole improvement results in significantly larger coefficients and, as we will show, greatly reduced finite- a errors. Note that, unlike some improvement schemes, perturbative improvement of this sort does *not* introduce new parameters to be tuned numerically; here the additional couplings in the action are computed, before the simulation, using tadpole-improved perturbation theory.

Perturbatively improved actions have already proven very successful in simulations of heavy-quark mesons like the Υ . In the nonrelativistic NRQCD quark action, both relativistic effects and finite- a corrections are introduced through nonrenormalizable corrections to the basic action. The detailed simulation results presented in [3] agree well with experiment, and many depend crucially on these corrections. Tadpole improvement was essential to this success; test runs without tadpole improvement underestimated relativistic effects by as much as a factor of two.

The standard Wilson action for gluons has finite- a errors of $\mathcal{O}(a^2)$, which are plainly visible in the static-quark potential computed on a lattice with $a \approx 0.4$ fm (see Fig. 1a). To reduce these errors we use a corrected action [4] that includes




Figure 1. Static-quark potential computed on 6^4 lattices with $a \approx 0.4$ fm using the $\beta = 4.5$ Wilson action and the improved action with $\beta_{\text{pl}} = 6.8$, $\beta_{\text{rt}} = -0.562$, and $\beta_{\text{pg}} = -0.0844$.

two new terms in addition to the usual plaquette term:

$$\begin{aligned}
 S[U] = & \beta_{\text{pl}} \sum_{\text{pl}} \frac{1}{3} \text{Re Tr}(1 - U_{\text{pl}}) \\
 & + \beta_{\text{rt}} \sum_{\text{rt}} \frac{1}{3} \text{Re Tr}(1 - U_{\text{rt}}) \\
 & + \beta_{\text{pg}} \sum_{\text{pg}} \frac{1}{3} \text{Re Tr}(1 - U_{\text{pg}}), \quad (1)
 \end{aligned}$$

Here U_{pl} is the 1×1 plaquette operator, U_{rt} is the 1×2 rectangle operator, and U_{pg} is the six-link parallelogram operator (path $\mu, \nu, \rho, -\mu, -\nu, -\rho$ where μ, ν, ρ are all different directions). In our work, we take the plaquette coupling β_{pl} as an input and compute the other two using perturba-

tion theory. By tadpole improving the one-loop results in [4] we obtain:

$$\beta_{\text{rt}} = -\frac{\beta_{\text{pl}}}{20 u_0^2} (1 + 0.4805 \alpha_s) \quad (2)$$

$$\beta_{\text{pg}} = -\frac{\beta_{\text{pl}}}{u_0^2} 0.03325 \alpha_s. \quad (3)$$

As expected, tadpole improvement significantly improves the convergence of the β_{rt} expansion. Following [1], we use the measured expectation value of the plaquette to determine both the value of the mean link u_0 and the QCD coupling constant α_s used in these expressions:

$$u_0 = \left(\frac{1}{3} \text{Re Tr} \langle U_{\text{pl}} \rangle \right)^{1/4}, \quad (4)$$

$$\alpha_s = \frac{\frac{1}{3} \text{Re Tr}(1 - \langle U_{\text{pl}} \rangle)}{3.06839}. \quad (5)$$

The couplings β_{rt} and β_{pg} are determined self-consistently with u_0 and α_s for a given β_{pl} . As in NRQCD, there is no tuning here of the couplings for the correction terms; tadpole-improved perturbation theory determines them in terms of the single bare coupling β_{pl} .¹

The static-quark potential computed using the improved gluon action is shown in Fig. 1b. As in the Wilson case (Fig 1a), the lattice spacing is about 0.4 fm. The dashed line in these plots is the standard infrared parameterization for the potential, $V(r) = Kr - \pi/12r + c$, adjusted to fit the on-axis values of the potential. Off-axis points deviate from the fit by as much as 35% for the Wilson theory, indicating a significant failure of rotation invariance due to finite- a errors. By contrast, the deviations are only 2–4% for the improved theory — negligible for most low-energy applications. Tuning the couplings away from their one-loop perturbative values results in little further improvement, suggesting that $\mathcal{O}(a^4)$ effects are as important at this a as the $\mathcal{O}(\alpha_s^2 a^2)$ errors that have not been included.

To further check on our improved theory, we examined the spin-averaged spectrum of the ψ family of mesons using NRQCD for the c -quarks and our improved action at $\beta_{\text{pl}} = 6.8$ and 7.4. Because

¹Using identities from [2] we find that our action is positive semidefinite at least for $\beta_{\text{pl}} \geq 6.8$.

	β_{pl}	a	$(1P - 1S)/\sqrt{\sigma}$
Wilson	4.5	.41 fm	1.46 (5)
	5.7	.17 fm	0.92 (5)
Improved	6.8	.41 fm	0.99 (5)
	7.4	.25 fm	0.93 (5)

Table 1

Ratio of the charmonium $1P - 1S$ splitting to the square root of the string tension.

we were examining only the spin-averaged spectrum, we omitted all relativistic corrections from the NRQCD action, but kept the corrections for $\mathcal{O}(a, a^2)$ errors. The spectra, normalized to give the correct $1P-1S$ splitting, are shown in Fig. 2 together with experimental results (dashed lines) and simulation results obtained using the Wilson action for the gluons at smaller a 's [3]. All agree to within 30–40 MeV. Also the results scale well with the string tension σ in the improved theory, but very poorly in the Wilson theory (see Table 1).

Since coarse lattices have far fewer sites and much less critical-slowing-down, the cost to produce a statistically independent configuration should be much less on a coarser lattice. To examine this issue, we compared our results with those in [5] for the potential computed using the Wilson action at $\beta = 6$ (32^4 lattice with $a \approx 0.1$ fm). We rescaled the coordinates and potential from this other study to put them in the same units as our $\beta_{\text{pl}} = 6.8$ results, and examined the potentials at comparable distances. The results from both simulations are listed in Table 2. In both cases the potential is obtained from the time dependence of loop-like correlation functions for times equal to or larger than some T_{min} . Our results required 1.3×10^7 site updates, while the analysis on the fine lattice required 6.4×10^9 site updates. Since statistical errors (for $T_{\text{min}} = 0.4$ fm) are about 20 times smaller for the coarse lattice, we estimate that comparable errors with the fine lattice would require 197,000 times more site updates than we used on the coarse lattice.

Comparisons of the sort just discussed are necessarily crude, but the implications are clear. The




Figure 2. S , P , and D states of charmonium computed on lattices with: $a \approx 0.41$ fm (improved action, $\beta_{\text{pl}} = 6.8$); $a \approx 0.25$ fm (improved action, $\beta_{\text{pl}} = 7.4$); and $a \approx 0.17$ fm (Wilson action, $\beta = 5.7$).

r/a_c	$a = a_c \approx 0.41$		$a \approx 0.10$
	$T_{\text{min}} = .4$	$T_{\text{min}} = .8$	$T_{\text{min}} = .3$
1	0.871 (0)	0.887 (1)	
0.962			0.839 (11)
$\sqrt{2}$	1.373 (1)	1.384 (2)	
1.361			1.316 (25)
$\sqrt{3}$	1.718 (2)	1.742 (10)	
1.667			1.581 (54)
2	1.897 (2)	1.941 (10)	
1.924			1.842 (42)

Table 2

Comparison of the static-quark potential $aV(r)$ as computed on a coarse lattice (improved action, $\beta_{\text{pl}} = 6.8$, $a \approx 0.41$ fm), and on a fine lattice (Wilson action, $\beta = 6$, $a \approx 0.10$ fm).

shift to coarse lattices and perturbatively improved actions, with tadpole-improved couplings, makes possible the calculation of the QCD hadron spectrum to, say, 1% accuracy with orders of magnitude less computing power than is possible with current methods.

We are very grateful to A. Pierce, IBM, and Cornell's Center for Theory and Simulation for help in using the IBM SP-2 supercomputer for parts of this work. This work was supported by the DOE and NSF.

REFERENCES

- [1] G.P. Lepage and P.B. Mackenzie, *Phys. Rev.* **D48**, 2250 (1993).
- [2] M. Lüscher and P. Weisz, *Comm. Math. Phys.* **97**, 59 (1985); K. Symanzik, *Nucl. Phys.* **B226**, 187(1983).
- [3] C. Davies et al, to be published in *Phys. Rev.* **D** (1994); and talks by C. Davies and J. Sloan at this meeting.
- [4] See M. Lüscher and P. Weisz, *Phys. Lett.* **158B**, 250(1985), and references therein.
- [5] G.S. Bali and K. Schilling, *Phys. Rev.* **D46**, 2636(1992).